

Supersymmetric Nonlinear Sigma Models on Ricci-flat Kähler Manifolds with $O(N)$ Symmetry

Kiyoshi Higashijima^{*}, Tetsuji Kimura[†] and Muneto Nitta[‡]

*Department of Physics, Graduate School of Science, Osaka University,
Toyonaka, Osaka 560-0043, Japan*

Abstract

We propose a class of $\mathcal{N} = 2$ supersymmetric nonlinear sigma models on the Ricci-flat Kähler manifolds with $O(N)$ symmetry.

^{*} E-mail: higashij@phys.sci.osaka-u.ac.jp

[†] E-mail: t-kimura@het.phys.sci.osaka-u.ac.jp

[‡] E-mail: nitta@het.phys.sci.osaka-u.ac.jp

String theory propagating in a curved spacetime is described by a conformally invariant nonlinear sigma model in two dimensions. Spacetime supersymmetry of the string theory requires $\mathcal{N} = 2$ worldsheet supersymmetry. The conformal invariance is realized in finite field theories with vanishing β -functions. In this letter, we propose a class of nonlinear sigma models on Ricci-flat Kähler manifolds with $O(N)$ symmetry.

Two-dimensional $\mathcal{N} = 2$ supersymmetric nonlinear sigma models are described by (anti-)chiral superfields: $\phi = A + \sqrt{2}\theta\psi + \theta\theta F$ ($\phi^\dagger = A^* + \sqrt{2}\bar{\theta}\bar{\psi} + \bar{\theta}\bar{\theta}F^*$) where A , ψ and F are complex scalar, Dirac fermion, auxiliary scalar fields, respectively. To define supersymmetric nonlinear sigma models on the Ricci-flat Kähler manifolds with $O(N)$ symmetry, we prepare dynamical chiral superfields ϕ^i ($i = 1, \dots, N$; $N \geq 2$), belonging to the vector representation of $O(N)$, and an auxiliary chiral superfield ϕ_0 , belonging to an $O(N)$ singlet. The most general Lagrangian with $O(N)$ symmetry, composed of these chiral superfields, is given by

$$\mathcal{L} = \int d^4\theta \mathcal{K}(x) + \int d^2\theta \phi_0 \left(\sum_{i=1}^N \phi^i \phi^i - a^2 \right) + \int d^2\bar{\theta} \phi_0^\dagger \left(\sum_{i=1}^N \phi^{\dagger i} \phi^{\dagger i} - a^2 \right), \quad (1)$$

where x is the $O(N)$ invariant defined by

$$x = \sum_{i=1}^N \phi^{\dagger i} \phi^i, \quad (2)$$

and $\mathcal{K}(x)$ is an arbitrary function of x . In the Lagrangian (1), we can assume that a^2 is a positive real constant, using the field redefinition. By the integration over the auxiliary field ϕ_0 , we obtain the constraint among the superfields ϕ^i , $\sum_{i=1}^N \phi^i \phi^i = a^2$, whose bosonic part is

$$\sum_{i=1}^N A^i A^i = a^2. \quad (3)$$

The manifold defined by this constraint with the Kähler potential $\mathcal{K}$ is a *non-compact* Kähler manifold with the complex dimension $N - 1$, where the $O(N)$ symmetry acts as a holomorphic isometry. One of the component, say A^N , can be expressed in terms of the independent fields A^a ($a = 1, \dots, N - 1$)

$$A^N = \sqrt{a^2 - \sum_{a=1}^{N-1} A^a A^a}. \quad (4)$$

To obtain the Ricci-flat Kähler manifold with $O(N)$ symmetry, we calculate the Ricci form and solve the Ricci-flat condition. To do this, we first calculate the Kähler metric $g_{ab} = \partial_a \partial_{\bar{b}} \mathcal{K}$ of the

manifold:

$$g_{ab^*} = \mathcal{K}' \left(\delta_{ab} + \frac{A^a A^{*b}}{A^{*N} A^N} \right) + \mathcal{K}'' \left(A^{*a} - \frac{A^{*N}}{A^N} A^a \right) \left(A^b - \frac{A^N}{A^{*N}} A^{*b} \right), \quad (5)$$

where the prime denotes differentiation with respect to x , and A^N is given by (4). Here we have used the same letter x for its lowest component: $x = \sum_{i=1}^N A^{*i} A^i$. Using this metric the nonlinear sigma model Lagrangian is written as

$$\mathcal{L} = g_{ab^*} \partial_\mu A^a \partial^\mu A^{*b} + i g_{ab^*} \bar{\psi}^b (\not{D} \psi)^a + \frac{1}{4} R_{ab^* cd^*} \psi^a \psi^c \bar{\psi}^b \bar{\psi}^d, \quad (6)$$

where $R_{ab^* cd^*}$ is the Riemann curvature tensor and $(D_\mu \psi)^a = \partial_\mu \psi^a + \partial_\mu A^b \Gamma_{bc}^a \psi^c$ [1]. Without loss of generality, an arbitrary point on the manifold can be transformed by the $O(N)$ symmetry to

$$A^\alpha = 0 \quad (\alpha = 1, 2, \dots, N-2), \quad A^{N-1} = z, \quad A^N = \sqrt{a^2 - z^2}, \quad (7)$$

where z is complex. The Kähler metric on this point is

$$g_{ab^*} = \begin{pmatrix} \mathcal{K}' \delta_{\alpha\beta^*} & 0 \\ 0 & \frac{1}{A^{*N} A^N} [\mathcal{K}' \cdot x + \mathcal{K}'' \cdot (x^2 - a^4)] \end{pmatrix}, \quad (8)$$

and its determinant is given by

$$\det g_{ab^*} = \frac{1}{A^{*N} A^N} [(\mathcal{K}')^{N-1} \cdot x + (\mathcal{K}')^{N-2} (\mathcal{K}'') \cdot (x^2 - a^4)]. \quad (9)$$

The Ricci form is given by $R_{ab^*} = -\partial_a \partial_{b^*} \log \det g$. Hence, the Ricci-flat condition $R_{ab^*} = 0$ is equivalent to the condition that the determinant is a constant up to products of holomorphic and anti-holomorphic functions: $\det g_{ab^*} \equiv (\text{constant}) \times F(z) \times F^*(z^*)$. From this condition, we obtain a differential equation

$$(\mathcal{K}')^{N-1} \cdot x + (\mathcal{K}')^{N-2} (\mathcal{K}'') \cdot (x^2 - a^4) = c, \quad (10)$$

where c is a constant. Using this equation, the metric (5) can be rewritten as

$$g_{ab^*} = \frac{\mathcal{K}'}{A^{*N} A^N} \left[\delta_{ab} A^{*N} A^N + A^a A^{*b} + \frac{c \cdot (\mathcal{K}')^{-(N-1)} - x}{x^2 - a^4} (A^{*a} A^N - A^a A^{*N}) (A^b A^{*N} - A^{*b} A^N) \right], \quad (11)$$

where A^N is given by (4). Therefore, we only need the solution of $\mathcal{K}'$ but not $\mathcal{K}$ itself, to calculate the Ricci-flat Kähler metric.

To solve the nonlinear differential equation (10), we transform it to a linear differential equation:

$$\frac{dy(x)}{dx} + (N-1)\frac{x}{x^2-a^4}y(x) = \frac{c(N-1)}{x^2-a^4}, \quad y(x) = (\mathcal{K}')^{N-1}. \quad (12)$$

The general solution of $y(x)$ is immediately obtained as

$$y(x) = \left(\frac{1}{x^2-a^4}\right)^{\frac{N-1}{2}} \left\{ y_0(x_0^2-a^4)^{\frac{N-1}{2}} + c(N-1) \int_{x_0}^x dt (t^2-a^4)^{\frac{N-3}{2}} \right\}, \quad (13)$$

where x_0 is an integration constant and $y_0 = y(x_0)$. In order to obtain the finite solution at $x = a^2$, we must set the parameter x_0 the solution of

$$y_0(x_0^2-a^4)^{\frac{N-1}{2}} + c(N-1) \int_{x_0}^{a^2} dt (t^2-a^4)^{\frac{N-3}{2}} = 0. \quad (14)$$

Using this equation, we obtain an integral representation of $y(x)$:

$$y(x) = (\mathcal{K}')^{N-1} = c(N-1) \left(\frac{1}{x^2-a^4}\right)^{\frac{N-1}{2}} \int_{a^2}^x dt (t^2-a^4)^{\frac{N-3}{2}}. \quad (15)$$

By performing the integration we can express $y(x)$ using the hypergeometric function $F(\alpha, \beta, \gamma; z)$:

$$\begin{aligned} (\mathcal{K}')^{N-1} &= \frac{c(N-1)}{2\Gamma(N/2)} (-a^4)^{\frac{N-3}{2}} \left(\frac{1}{x^2-a^4}\right)^{\frac{N-1}{2}} \\ &\times \left[2x\Gamma\left(\frac{N}{2}\right) \cdot F\left(\frac{1}{2}, -\frac{N-3}{2}, \frac{3}{2}; \frac{x^2}{a^4}\right) - a^2\sqrt{\pi}\Gamma\left(\frac{N-1}{2}\right) \right]. \end{aligned} \quad (16)$$

We can also obtain the Ricci-flat metric by substituting $\mathcal{K}'$ to (11). We thus have obtained $(N-1)$ -dimensional Ricci-flat Kähler manifolds with $O(N)$ symmetry.

When N is odd ($N = 2m+1$ and $m = 1, 2, \dots$), the hypergeometric function reduces to a polynomial:

$$(\mathcal{K}')^{2m} = 2c \sum_{n=0}^{m-1} {}_{m-1}C_n \frac{m}{2n+1} (-a^4)^{m-n-1} \left[\frac{x^{2n+1} - (a^2)^{2n+1}}{(x^2-a^4)^m} \right], \quad {}_mC_n = \frac{m!}{n!(m-n)!}. \quad (17)$$

If N is even ($N = 2m+2$ and $m = 1, 2, \dots$) except for $N = 2$ [see (19a), below, for the $N = 2$ solution], the solution can be written as

$$\begin{aligned} (\mathcal{K}')^{2m+1} &= \frac{c(2m+1)}{a^2(\tan \theta_x)^{2m+1}} \sum_{n=0}^m {}_mC_n (-1)^{m-n} \\ &\times \frac{(2n-1)!!}{(2n)!!} \left\{ \frac{\sin \theta_x}{(\cos \theta_x)^{2n}} \sum_{r=0}^{n-1} \frac{(2n-2r-2)!!}{(2n-2r-1)!!} (\cos \theta_x)^{2r} + \log \left| \tan \left(\frac{\theta_x}{2} + \frac{\pi}{4} \right) \right| \right\}, \end{aligned} \quad (18)$$

where $\theta_x = \arccos(a^2/x)$. Explicit expressions of $\mathcal{K}'$ for $N = 2$ to 7 are

$$N = 2 : \mathcal{K}' = \frac{c}{\sqrt{x^2-a^4}} \log \left(\frac{x + \sqrt{x^2-a^4}}{a^2} \right), \quad (19a)$$

$$N = 3 : (\mathcal{K}')^2 = \frac{2c}{x+a^2}, \quad (19b)$$

$$N = 4 : (\mathcal{K}')^3 = \frac{3c}{2} \frac{1}{(x^2-a^4)^{3/2}} \left[x\sqrt{x^2-a^4} - a^4 \log \left(\frac{x+\sqrt{x^2-a^4}}{a^2} \right) \right], \quad (19c)$$

$$N = 5 : (\mathcal{K}')^4 = \frac{4c}{3} \frac{x+2a^2}{(x+a^2)^2}, \quad (19d)$$

$$N = 6 : (\mathcal{K}')^5 = -\frac{5c}{8} \frac{1}{(x^2-a^4)^{5/2}} \left[(-2x^3+5a^4x)\sqrt{x^2-a^4} - 3a^8 \log \left(\frac{x+\sqrt{x^2-a^4}}{a^2} \right) \right], \quad (19e)$$

$$N = 7 : (\mathcal{K}')^6 = \frac{2c}{5} \frac{3x^2+9a^2x+8a^4}{(x+a^2)^3}. \quad (19f)$$

For definiteness, we give the explicit solution of $\mathcal{K}$ for $N = 2$ and $N = 3$. For $N = 2$, $\mathcal{K}$ can be obtained as

$$\mathcal{K}_{N=2} = \frac{c}{2} \left[\log \left(\frac{x+\sqrt{x^2-a^4}}{a^2} \right) \right]^2. \quad (20)$$

By the field redefinition $A^1 = a \cos \varphi$ ($\varphi \in \mathbf{C}$), this Kähler potential becomes

$$\mathcal{K}_{N=2} = c\varphi^*\varphi + F(\varphi) + F^*(\varphi^*). \quad (21)$$

Here, $F(\varphi)$ and $F^*(\varphi^*)$ are holomorphic and anti-holomorphic functions, respectively, which can be eliminated by the Kähler transformation. Thus we obtain the free field theory with the flat metric $g_{\varphi\varphi^*} = c$ as we expected, since the Ricci-flatness implies the vanishing Riemann curvature in real two-dimensional manifolds. The solution of $\mathcal{K}$ for $N = 3$ can be calculated, to yield

$$\mathcal{K}_{N=3} = \sqrt{8c} \sqrt{x+a^2}. \quad (22)$$

The Ricci-flat Kähler metric for $N = 3$ is

$$g_{ab^*} = \frac{\sqrt{2c}}{(x+a^2)^{3/2}} \left[(x+a^2) \left(\delta_{ab} + \frac{A^a A^{*b}}{A^{*3} A^3} \right) - \frac{1}{2} \left(A^{*a} - \frac{A^{*3}}{A^3} A^a \right) \left(A^b - \frac{A^3}{A^{*3}} A^{*b} \right) \right], \quad (23)$$

where A^3 is given in (4). This defines a (real) four-dimensional hyper-Kähler manifold with $O(3)$ symmetry, the Eguchi-Hanson space [2]. The metric for $N = 4$ calculated from (19c) coincides with the metric of the deformed conifold, obtained earlier in [3].

We discuss the limit of vanishing a^2 . In this limit, the manifold becomes a conifold, in which the point represented by $z = 0$ in (7) is singular. In this limit, we can obtain the explicit solution of $\mathcal{K}$ for any $N(> 2)$, given by

$$\mathcal{K} = \left(c \frac{N-1}{N-2} \right)^{\frac{1}{N-1}} \int_{x_0}^x dt t^{\frac{-1}{N-1}} = \frac{1}{c} \left(c \frac{N-1}{N-2} \right)^{\frac{N}{N-1}} \cdot x^{\frac{N-2}{N-1}} + (\text{constant}). \quad (24)$$

In the $N \rightarrow \infty$ limit, the Kähler potential (24) becomes the simplest form:

$$\lim_{N \rightarrow \infty} \mathcal{K} = x . \quad (25)$$

Therefore, when we discuss non-perturbative effects of the sigma model on the conifold defined by (24), using the $1/N$ expansion method, it is sufficient to consider the simplest Kähler potential $\mathcal{K} = x$ instead of (24).

If we prepare invariants of other groups in the superpotential of (1), we would obtain Ricci-flat Kähler manifolds with other symmetry.

We have studied the nonlinear sigma models on Ricci-flat Kähler manifolds. These models have vanishing β -function up to the fourth order in the perturbation theory [4, 5]. Despite the appearance of non-zero β -function at the four-loop order, we will be able to obtain the conformally invariant field theory for the background metric related to the Ricci-flat manifolds through non-local field redefinition [6].

After the completion of this work we came to know that our metric was also discussed in other context [7, 8]. Let us give a comment on the relation to their work. Defining the new variable u by the relation $(a^2 \sinh 2u)^2 = t^2 - a^4$ in Eq. (15), the integral can be rewritten as $2a^{2(N-2)} \int_0^r (\sinh 2u)^{N-2} du = 2a^{2(N-2)} R(r)$, where r has been defined by $(a^2 \sinh 2r)^2 = x^2 - a^4$. Here r and $R(r)$ are the same ones introduced in [8]. We obtain the Kähler potential differentiated by r , given by

$$\frac{d\mathcal{K}}{dr} = 2a^2 \left\{ \frac{2c}{a^2} (N-1) \right\}^{\frac{1}{N-1}} R(r)^{\frac{1}{N-1}} . \quad (26)$$

Acknowledgements

We would like to thank Gary Gibbons for pointing out references [8]. We are also grateful to Takashi Yokono for useful comments.

References

- [1] B. Zumino, Phys. Lett. **87B** (1979) 203.
- [2] T. Eguchi and A. J. Hanson, Phys. Lett. **74B** (1978) 249.

- [3] P. Candelas and X. C. de la Ossa, Nucl. Phys. **B342** (1990) 246.
K. Ohta and T. Yokono, JHEP **0002** (2000) 023, [hep-th/9912266](#).
- [4] L. Alvarez-Gaumé, Nucl. Phys. **B184** (1981) 180.
C. M. Hull, Nucl. Phys. **B260** (1985) 182.
L. Alvarez-Gaumé and P. Ginsparg, Comm. Math. Phys. **102** (1985) 311.
- [5] M. T. Grisaru, A. E. M. van de Ven and D. Zanon, Phys. Lett. **173B** (1986) 423; Nucl. Phys. **B277** (1986) 388; Nucl. Phys. **B277** (1986) 409.
- [6] D. Nemeschansky and A. Sen, Phys. Lett. **178B** (1986) 365.
- [7] M. B. Stenzel, Manuscripta Math. **80** (1993) 151.
- [8] M. Cvetič, G. W. Gibbons, H. Lü and C. N. Pope, [hep-th/0012011](#); [hep-th/0102185](#).